Author Index

Alencar, J. De, see Barreira, E.S. 381
Alessi, D. M., see Pirkle, W. H. 203
Arancibia, E. L.
——, Castells, R. C. and Nardillo, A. M.
Thermodynamic study of the behaviour of two molten organic salts as stationary phases in gas chromatography 21
Arpino, P. J., see Cousin, J. 125
Barefoot, A. C.
—— and Reiser, R. W.
Packed capillary liquid chromatography–mass spectrometry using both direct-coupling and moving-belt interfaces 217
Barnes, S., see Hayes, S. M. 239
Barreira, E. S.
——, Parente, J. P. and De Alencar, J. W.
Separation and determination of aliphatic amines by high-performance liquid chromatography with ultraviolet detection 381
Bayer, H., see Klaus, R. 300
Baziotis, A., see Whithka, J. 175
Beger, J., see Voelkel, A. 31
Bender, P. E., see Erhard, K. F. 360
Berezkin, V. G., see Viktorova, E. N. 317
Berkowitz, S. A.
Manipulation of the mobile phase to achieve multiple step protein (calmodulin) purification using the same chromatographic material (a weak anion exchanger) 288
Berni, R. J., see Hayes, S. M. 239
Bezemer, R., see Van den Oetelaar, P. J. M. 323
Blake, J. D.
——, Clarke, M. L. and Richards, G. N.
Determination of organic acids in sugar cane process juice by high-performance liquid chromatography. Improved resolution using dual Aminex HPX-87H cation-exchange columns equilibrated to different temperatures 265
Borák, J., see Konečný, P. 387
Bousfield, G. R., see Ward, D. N. 255
Brieva, A., see Rubio, N. 366
Castagnoli, N., Jr., see Shinka, T. 279
Castells, R. C., see Arancibia, E. L. 21
Chadha, R. K., see Lawrence, J. F. 355
Charbonneau, G. M., see Naikwadi, K. P. 227
Chen, T.-M.
——, Contario, J. J. and Fike, R. R.
High-performance liquid chromatographic assay for vigabatrin and its primary degradation product in a pharmaceutical tablet formulation 351
Choi, C. K., see Kim, H. S. 372
Clarke, M. Ł., see Blake, J. D. 265
Clement, R. E., see Naikwadi, K. P. 227
Colmsjö, A. L.
—— and Ericsson, M. W.
Assessment of the height equivalent to a theoretical plate in liquid chromatography 63
Contario, J. J., see Chen, T.-M. 351
Cossu, G.
—— and Righetti, P. G.
Resolution of G_γ and A_γ foetal haemoglobin tetramers in immobilized pH gradients 211
Cousin, J.
—— and Arpino, P. J.
Construction of a supercritical fluid chromatograph–mass spectrometer instrument system using capillary columns, and a chemical ionization source accepting high flow-rates of mobile phase 125
De Alencar, J. W., see Barreira, E. S. 381
Ebert, K., see Voelkel, A. 31
Erhard, K. F.
——, Razgaitis, K. A. and Bender, P. E.
Purification of leukotriene B_4 by semi-preparative high-performance liquid chromatography 360
Ericsson, M. W., see Colmsjö, A. L. 63
Faurie, M., see Foucault, A. 294
Fike, R. R., see Chen, T.-M. 351
Fischer, W., see Klaus, R. 300
Foucault, A.
—— and Faurie, M.
Ion-exchange separation and pK_a determinations of the α- and β-isomers of N-acetyl-L-aspartyl-L-glutamic acid. Correlation of chromatographic properties with pK_a values 294
Fox, T. R., see Kastl, P. E. 347
García-Raso, A.
——, Martínez-Castro, I., Páez, M. I., Sanz, J., García-Raso, J. and Sauro-Calixto, F.
Gas chromatographic behaviour of carbohydrate trimethylsilyl ethers. I. Aldopentoses 9
García-Raso, J., see García-Raso, A. 9
Gavrichev, V. S., see Viktorova, E. N. 317
Gill, R., see Smith, R. M. 73
Gioeli, C., see Pettersson, C. 247
Hansen, B., see Welinder, B. S. 309
Hashimoto, T., see Kato, Y. 327, 340

Hayes, S. M.
——, Liu, R. H., Tsang, W.-S., Legendre, M. G., Berni, R. J., Pillion, D. J., Barnes, S. and Ho, M. H.
Enantiomeric composition analysis of amphetamine and methamphetamine by chiral phase high-performance liquid chromatography–mass spectrometry 239
Hermann, E. A., see Kastl, P. E. 347
Herraiz, M., see Loyola, E. 53
Ho, M. H., see Hayes, S. M. 239
Hoag, M. K. P., see Shinka, T. 279
Hoenders, H. J., see Van den Oetelaar, P. J. M. 323
Horváth, Cs., see Kalghatgi, K. 335
Hurdley, T. G., see Smith, R. M. 73
Hyun, M. H., see Pirkle, W. H. 203
Ikeda, H., see Tanaka, M. 165
Kalghatgi, K.
—— and Horváth, Cs.
Rapid analysis of proteins and peptides by reversed-phase chromatography 335
Karasek, F. W., see Naikwadi, K. P. 227
Kastl, P. E.
——, Fox, T. R. and Hermann, E. A.
Quantitative determination of 1,2-butylene oxide in whole rat blood by gas chromatography–mass spectrometry 347
Kato, Y.
——, Kitamura, T., Mitsui, A. and Hashimoto, T.
High-performance ion-exchange chromatography of proteins on non-porous ion exchangers 327
——, Nakamura, K. and Hashimoto, T.
High-performance hydroxyapatite chromatography of proteins 340
Kim, H. S.
——, Choi, C. K. and Park, Y. H.
Determination of cholesterol and its fermentation products by high-performance liquid chromatography 372
Kitamura, T., see Kato, Y. 327
Klaus, R.
——, Fischer, W. and Bayer, H.
Praktische Erfahrungen über Nachweis und dünnschichtchromatographische Trennung von Phenolen im Hinblick auf eine quantitative Analyse 300
Klesper, E., see Leyendecker, D. 89, 105
Konečný, P.
——, Smrž, M., Borák, J. and Slováková, S.
Affinity chromatography of pig heart malate dehydrogenase on the Remazol Brilliant Blue R derivative of hydroxyethyl methacrylate gel 387
Korhonen, I. O. O., see Kuchař, M. 43
Kuchař, M.
——, Tomková, H., Rejholec, V. and Korhonen, I. O. O.
Gas–liquid chromatography and lipophilicity of esters of benzoic acids. Enthalpy–entropy compensation 43
Kunugi, A.
—— and Tabei, K.
On-column organic reactions. Schotten-Baumann benzoylation of phenols on Extrelut column 320
Lawrence, J. F.
—— and Chadha, R. K.
Headspace liquid chromatographic technique for the determination of sulfite in food 355
Legendre, M. G., see Hayes, S. M. 239
Lehtonen, P.
Molecular connectivity indices in the prediction of the retention of oxygen-containing amines in reversed-phase liquid chromatography 143
Leyendecker, D.
——, Leyendecker, D., Lorenschat, B., Schmitz, F. P. and Klesper, E.
Influence of density on the chromatographic behaviour of lower alkanes as mobile phases in supercritical fluid chromatography 89
——, Leyendecker, D., Schmitz, F. P., Lorenschat, B. and Klesper, E.
Supercritical fluid chromatography using mixtures of carbon dioxide or ethane with 1,4-dioxane as eluents 105
Liu, R. H., see Hayes, S. M. 239
Lorenschat, B., see Leyendecker, D. 89, 105
Loyola, E.
——, Herraiz, M., Reglero, G. and Martin-Alvarez, P.
Evaluation of a programmed temperature vaporizer for quantitative analysis by split injection 53
Martin-Alvarez, P., see Loyola, E. 53
Martínez-Castro, I., see García-Raso, A. 9
Maskiewicz, R., see Whithka, J. 175
Matsumoto, U., see Shibusawa, Y. 153
Mitsui, A., see Kato, Y. 327
Moncuse, P., see Whithka, J. 175
Naikwadi, K. P.
——, Charbonneau, G. M., Karasek, F. W. and Clement, R. E.
Separation and indentification of organic compounds in air particulate extracts by high-performance liquid chromatography and gas chromatography–mass spectrometry 227
Nakamura, K., see Kato, Y. 340
Nardillo, A. M., see Arancibia, E. L. 21
Oetelaar, P. J. M. van den, see Van den Oetelaar, P. J. M. 323

Osselton, M. D., see Smith, R. M. 73
Páez, M. I., see García-Raso, A. 9
Parente, J. P., see Barreira, E. S. 381
Park, Y. H., see Kim, H. S. 372
Pettersson, C.
—— and Gioeli, C.
Separation of enantiomeric acids using immobilized acetylquinine as a chiral stationary phase 247
Pilc, J. A.
—— and Sermon, P. A.
Chromatographic separation of a wide range of ethoxylated non-ionic surfactants and their sulphonates using high-performance liquid chromatography 375
Pillion, D. J., see Hayes, S. M. 239
Pirkle, W. H.
——, Alessi, D. M., Hyun, M. H. and Pochapsky, T. C.
Separation of some enantiomeric di- and tripeptides on chiral stationary phases 203
Pochapsky, T. C., see Pirkle, W. H. 203
Razgaitis, K. A., see Erhard, K. F. 360
Reglero, G., see Loyola, E. 53
Reiser, R. W., see Barefoot, A. C. 217
Rejholec, V., see Kuchař, M. 43
Richards, G. N., see Blake, J. D. 265
Righetti, P. G., see Cossu, G. 211
Rubio, N.
—— and Brieva, A.
Purification of allergens by high-performance liquid chromatography. III. Purification of fungal allergens 366
Sanz, J., see García-Raso, A. 9
Saura-Calixto, F., see García-Raso, A. 9
Schmitz, F. P., see Leyendecker, D. 89, 105
Sermon, P. A., see Pilc, J. A. 375
Shibusawa, Y.
——, Matsumoto, U. and Takatori, M.
Surface affinity chromatographic separation of blood cells. VI. Capacity factors of human peripheral blood cells on poly(propylene glycol) bonded Chromagel columns and their correlation with surface hydrophobicities of the gel beads and the cells 153
Shinka, T.
——, Castagnoli, N., Jr., Wu, E. Y., Hoag, M. K. P. and Trevor, A. J.
Cation-exchange high-performance liquid chromatography assay for the nigrostriatal toxicant 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine and its monoamine oxidase B generated metabolites in brain tissues 279
Shono, T., see Tanaka, M. 165
Slováková, S., see Konečný, P. 387
Smith, R. M.
——, Hurdley, T. G., Gill, R. and Osselton, M. D.
Retention reproducibility of basic drugs in high-performance liquid chromatography on a silica column with a methanol–ammonium nitrate eluent. The effect of the mobile phase and the operating conditions 73
Smrž, M., see Konečný, P. 387
Sørensen, H. H., see Welinder, B. S. 309
Szymanowski, J., see Voelkel, A. 31
Tabei, K., see Kunugi, A. 320
Takatori, M., see Shibusawa, Y. 153
Tanaka, M.
——, Ikeda, H. and Shono, T.
Carbamoylated cyclodextrin-bonded stationary phases for liquid chromatography 165
Tomková, H., see Kuchař, M. 43
Trevor, A. J., see Shinka, T. 279
Tsang, W.-S., see Hayes, S. M. 239
Van den Oetelaar, P. J. M.
——, Bezemer, R. and Hoenders, H. J.
Fast protein liquid chromatography–chromatofocusing of bovine and chicken α-crystallin subunits under denaturing conditions 323
Viktorova, E. N.
——, Berezkin, V. G. and Gavrichev, V. S.
Gas chromatography of alkenes in water vapour as carrier gas on a solid support impregnated with silver nitrate 317
Voelkel, A.
——, Szymanowski, J., Beger, J. and Ebert, K.
Polarity of 1,3-bis[ω-butoxyoligo(oxyethylene)]propan-2-ols as measured by reversed-phase gas chromatography 31
Ward, D. N.
——, Wen, T. and Bousfield, G. R.
Sulfate and phosphate analysis in glycoproteins and other biologic compounds using ion chromatography. Application to glycoprotein hormones and sugar esters 255
Welinder, B. S.
——, Sørensen, H. H. and Hansen, B.
Reversed-phase high-performance liquid chromatography of human growth hormone 309
Wen, T., see Ward, D. N. 255
Whithka, J.
——, Moncuse, P., Baziotis, A. and Maskiewicz, R.
Use of high-performance size-exclusion, ion-exchange, and hydrophobic interaction chromatography for the measurement of protein conformational change and stability 175
Wu, E. Y., see Shinka, T. 279